



Li StarFish S.r.l.
Via Cavour, 35 - 20063 Cernusco S/N (MI), Italy
Tel. +39-02-92150794 - Fax. +39-02-92157285
info@listarfish.it - www.listarfish.it

ELISA for Mouse IgG

Product Code: 3825-1AD-6

CONTENTS:

Vial 1 (yellow top)

Anti-IgG antibody (150 μ l)

Concentration: 0.5 mg/ml

Vial 2 (green top)

ALP-conjugated anti-IgG antibody (80 μ l)

Vial 3

Lyophilised mouse IgG standard

To ensure total recovery of stated quantity, vials have been overfilled.

STORAGE:

Shipped at ambient temperature. On arrival box 1 should be stored refrigerated at 4-8°C and box 2 should be stored frozen at -20°C.

General

Intended use: For quantitative determination of native mouse IgG in serum and plasma.

Reagents: Anti-IgG antibody is supplied in sterile-filtered (0.2 μm) PBS with sodium azide (0.02%). ALP-conjugated anti-IgG antibody is supplied in 0.1 M Tris-buffer with sodium azide (0.02%).

Recommended standard dilution: 0.1-500 ng/ml

Guidelines for Mouse IgG ELISA

- Day 1**
1. Coat a high protein binding ELISA plate with anti-IgG antibody, diluted to 1 $\mu\text{g/ml}$ in PBS, pH 7.4, by adding 100 $\mu\text{l/well}$. Incubate overnight at 4-8°C.
- Day 2**
2. Wash twice with PBS (200 $\mu\text{l/well}$).
 3. Block plate by adding 200 $\mu\text{l/well}$ of PBS with 0.05% Tween 20 (PBS-Tween) containing 0.1% BSA (incubation buffer*). Incubate for 1 hour at room temperature.
 4. Wash five times with PBS-Tween.
 5. Prepare mouse IgG standard by reconstituting contents of vial 3 in 500 μl PBS to a make up a stock solution of 50 $\mu\text{g/ml}$. The stock solution should be used immediately or stored in aliquots at -20°C for future use. We recommend the aliquots not be refrozen after initial use. For the test, prepare dilutions of the stock using the standard range as a guideline.
 6. Add 100 $\mu\text{l/well}$ of samples or standards diluted in incubation buffer and incubate for 2 hours at room temperature.
 7. Wash as in step 4.
 8. Add 100 $\mu\text{l/well}$ of anti-IgG-ALP diluted 1:1000 in incubation buffer. Incubate for 1 hour at room temperature.
 9. Wash as in step 4.
 10. Add 100 $\mu\text{l/well}$ of appropriate substrate solution e.g. p-nitrophenyl-phosphate (pNPP).
 11. Measure the optical density (405 nm for pNPP) in an ELISA reader after suitable developing time.

* The same buffer is used for blocking and for dilution.

NOTE; for research use only.

Li StarFish distribuisce:

